

# Contralateral Extremity Paresis and Numbness Caused by Pos-tero lateral Disc Herniation at C3–C4 in a Teenager

Fengbin Yu<sup>1</sup>, Fei Huang<sup>1</sup>, Hui Zhu<sup>1</sup>, Degang Tao<sup>1</sup>, Jianbo Jin<sup>1</sup>, and Lian Cen<sup>2\*</sup>

<sup>1</sup>Department of Orthopaedic Surgery, No. 72 Group Army Hospital of PLA, China

<sup>2</sup>Department of Product Engineering, School of Chemical Engineering, East China University of Science and Technology, China

Volume 1 Issue 3- 2018

Received Date: 11 Oct 2018

Accepted Date: 26 Oct 2018

Published Date: 02 Nov 2018

## 1. Abstract

We report a rare case of a teenager presenting with progressive contralateral extremity paresis and numbness caused by posterolateral disc herniation at C3-C4. Intervertebral decompression and artificial disc replacement was performed. Follow-up at 2 months showed complete neurologic recovery. The increasing tensile stresses from the dentate ligament during flexion of the neck might be the etiology of myelopathy in this patient.

## 2. Introduction

Cervical disc herniation at C3–C4 accounts for an estimated 4% of all cervical disc herniations [1] and is very rare in teenagers. The presenting symptoms often appear as myelopathy, Brown-Sequard syndrome, and seldom as radiculopathy in the few studies found in the literature [2-6]. We report a rare case of C3–C4 disc herniation causing a mild left compression of the spinal cord resulting in right extremity paresis and numbness.

## 3. Case Report

An 18-year-old boy presented with progressive paresis and numbness of the right upper and lower extremity of 3 weeks' duration. Lateral radiographs revealed a greater than normal flexion of the neck but no instability. MRI revealed a moderate left posterolateral disc herniation at C3–C4 with mild compression of the spinal cord (Figure 1). High signal intensity with a length of approximately 2 cm was visible on T2-weighted MRI (Figure 2). Conservative treatment was instituted with a restraining collar limiting mobility of the cervical spine but no improvement was seen. Progressive neurologic deterioration resulted in gait disturbance and right hand clumsiness. Neurological examination revealed reduced sensation to pain and temperature below the C4 dermatome and a decrease (MRC Grade 3/5) in motor function in the right extremity. Physical examination revealed positive Hoffman and Babinski signs in the right limb. Because of progressive neurologic deterioration and ineffective conservative treatment, intervertebral decompression and artificial disc replacement were performed. Follow-up at 2 months revealed normal motor strength and sensation in the right extremity.

Postoperative radiographs demonstrated good realignment of the cervical spine (Figure 3).




Figure 1: MRI revealed a moderate left posterolateral disc herniation at C3–C4 with mild compression of the spinal cord.




Figure 2: High signal intensity with a length of approximately 2 cm was visible on T2-weighted MRI.



Figure 3: Postoperative radiographs demonstrated good realignment of the cervical spine.

\*Corresponding Author (s): Lian Cen, Department of Product Engineering, School of Chemical Engineering, East China University of Science and Technology, Shanghai, 210000, China. E-mail: liancen1978@sina.com

Co-Corresponding author: Fei Huang, Department of Orthopaedic Surgery, No. 72 Group Army Hospital of PLA, Huzhou, 313000, China. E-mail: feihuang1965@sina.com

#### 4. Discussion

Cervical disc herniation of C3–C4 usually presents with quite different symptoms from the more commonly recognized lower cervical disc herniation [5]. As an unusual presentation of myelopathy, we report a rare case of C3–C4 left disc herniation resulting in right extremity dysfunction.

In this patient, it was very unusual that the mild compression of the spinal cord led to such severe neurological defects and that the left compression caused right extremity dysfunction. The theory of mechanical compression in which a herniated disc causes compression of the cord, leading to local tissue ischemia, injury, and neurological impairment certainly fails to explain the entire clinical presentation. Although the role of dentate ligaments in the etiology of myelopathy remains questionable [7-10], the theory of tensile stresses from the dentate ligaments appears to explain the clinical findings in this case well. First, the dentate ligament fibers run mediolaterally along the spinal cord with the dural attachments anchored by the dural root sleeves and dural ligaments and these are relatively fixed [11]. Tensile stresses transmitted to the spinal cord from the dura via the dentate ligaments may occur from a disc herniation as a result of displacing the spinal cord dorsally with less displacement of the dural attachments. Second, inferred from the clinical symptoms, the site of neurological injury was in the right corticospinal and left spinothalamic tracts. Both are located in the lateral columns of the spinal cord. Dentate ligament fibers also originate from the lateral columns. Therefore, the corticospinal and spinothalamic tracts should be most vulnerable because of lateral pulling by the dentate ligaments. Third, the arch of cervical flexion in this case was apparently larger than normal. Stretching on neural tissues by the dentate ligaments can be greatly amplified during a larger cervical flexion and can exceed the material properties of the tissue, leading to tissue disruption and transient or permanent neurological injury [12]. Last, the unbalanced tensile stresses transmitted to the spinal cord from the dura via the dentate ligaments generated by the non-centered disc herniation might be responsible for the asymmetric nerve damage seen in this case.

Removal of the herniated disc means removal of the tensile stresses transmitted to the spinal cord from the dura via the dentate ligaments, and early decompression through an anterior surgical approach is crucial to obtain good results.

**Acknowledgement:** This work was supported in part by 1112 Talent and Science technology Project of Huzhou (2017GY38) and New medical youngster project of Zhejiang

#### References

1. A Bucciero, L Vizioli, A Cerillo. Soft cervical disc herniation. An analysis of 187 cases. *J Neurosurg Sci.* 1998; 42: 125-130.
2. F T Sayer, A M Vitali, H L Low. Brown-Sèquard syndrome produced by C3-C4 cervical disc herniation: a case report and review of the literature. *Spine.* 2008; 33 (2008):PP. E279-282.
3. J T Kim, H J Bong, D S Chung. Cervical Disc Herniation Producing Acute Brown-Sequard Syndrome. *J Korean Neurosurg Soc.* 2009; 45: 312 - 314.
4. J Pan, L Li, L Qian. Intradural cervical disc herniation: report of two cases and review of the literature. *Spine.* 2001; 36: E1033-1037.
5. F T Sayer, A M Vitali, S Paquette. Isolated C3-C4 disc herniations present as a painless myelopathy. *Spine J.* 2008; 8: PP.729 - 731.
6. J K Lee, Y S Kim, S H Kim. Brown-Sequard syndrome produced by cervical disc herniation with complete neurologic recovery: report of three cases and review of the literature. *Spinal Cord.* 2007; 45: 744 - 748.
7. D N Levine. Pathogenesis of cervical spondylotic myelopathy. *J Neurol Neurosurg Psychiatry.* 1997; 62: 334 - 340.
8. S N Bishara. The posterior operation in treatment of cervical spondylosis with myelopathy: A long-term follow-up study. *J Neurol Neurosurg Psychiatry.* 1971; 34: PP. 393 - 398.
9. H F Stoltmann, W Blackwood. An anatomical study of the role of the dentate ligaments in the cervical spinal canal. *J Neurosurg.* 1966; 24: 43 - 46.
10. J F Cusick, J J Ackmann, S J Larson. Mechanical and physiological effects of dentotomy. *Neurosurg.* 1977; 46: PP. 767-775.
11. A F Tencer, B L Allen, R L Ferguson. A biomechanical study of thoracolumbar spinal fractures with bone in the canal: Part III, mechanical properties of the dura and its tethering ligaments. *Spine.* 1985; 10: 741-747.
12. F C Henderson, J F Geddes, A R Vaccaro. Stretch-associated injury in cervical spondylotic myelopathy: new concept and review. *Neurosurgery.* 2005; 56: 1101-1113.