

A Protruded Leision with Hemorrhage of Unexpected Histological Findings

Li Y, Wang Z and Yang J*

Department of Gastroenterology & Hepatology, West China Hospital of Sichuan University, Sichuan University–Oxford University Huaxi Gastrointestinal Cancer Centre, China

*Corresponding author:

Jinlin Yang,
Department of Gastroenterology and Hepatology,
West China Hospital, Sichuan University, 37# Guoxue
Lane, Chengdu, Sichuan, China. Postcode: 610041,
Tel: +86-18980602058, Fax: +86-28-85423387,
E-mail: yangjinlin@wchscu.cn

Received: 28 Sep 2021

Accepted: 13 Oct 2021

Published: 19 Oct 2021

Copyright:

©2021 Yang J. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

Citation:

Yang J, A Protruded Leision with Hemorrhage of Unexpected Histological Findings. *Ann Clin Med Case Rep.* 2021; V7(12): 1-3

Keywords:

Gastrointestinal bleeding; Renal cell carcinoma; Colonic protruded lesion; Direct invasion

Author contributions:

Yaru Li wrote the manuscript, Zhu Wang edited the manuscript. Jinlin Yang edited and revised the manuscript for intellectual content and is the article guarantor. All authors declared that neither the entire paper nor any part of its content has been published or has been accepted elsewhere. It is not being submitted to any other journal.

1. Clinical Image

An 83-year-old man complained of repeated hematochezia for 3 months. No other medical history was reported except for the percutaneous needle aspiration of left renal cyst 10 years ago. Colonoscopy revealed a 25-mm protruded lesion with central ulcer formation at the descending colon (Figure 1A). Massive bleeding occurred during forceps biopsy (Figure 1B), which was temporary managed by the clips. The initial pathologic result only indicated inflammation without evidence of malignancy. Abdominal enhanced computed tomography (CT) revealed a neoplastic lesion in the descending colon, involving adjacent left kidney (Figure 1C).

The laparoscopic surgery was indicated for the management of massive bleeding. However, the histopathological findings showed renal cell carcinoma (RCC) directly invaded from colonic serosa to propria muscularis with immunohistochemical staining expressing PAX-8, which indicated clear cell renal cell carcinoma (ccRCC)

(Figure 2A, H&E, orig. mag. $\times 40$; Figure 2B, H&E, orig. mag. $\times 200$; Figure 2C, IHC, orig. mag. $\times 200$). Renal cell carcinoma (RCC), located in the retroperitoneal cavity, usually metastasize to lymph nodes, lungs, bones, livers and brains [1], however, due to the anatomic separation, the metastasis from RCC to colon or primary colon tumor to kidney were only reported in few widespread cases, while direct invasion rarely occurred [2,3]. A case report observed that clips migrated to the descending colon after left kidney nephrectomy, possibly caused by the postoperative repairing process [4]. In the present case, the postoperative histological observation of the tumor revealed close adhesions between involved left kidney and colonic wall, which implied the rare site of colonic metastasis might occur as the result of the previous intervention of left renal cyst. Therefore, when a colonic lesion invaded nearby kidneys, the possibility of colonic metastasis of RCC needs to be considered, especially in the patients with history of renal disease or interventions.




Figure 1: **A:** Colonoscopy revealed a protruded lesion about 25 mm in diameter at the descending colon, occupying the lumen for 1/3 week with central ulcer formation; **B:** Spurting bleeding in the site of the biopsy; **C:** A mass-like tumor was observed in the descending colon and adjacent left kidney.




Figure 2: **A:** Histopathological findings showed clear cell renal cell carcinoma directly invaded from colonic serosa to propria muscularis (asterisk), the remaining mucosal and submucosal layers remained normal (H&E staining, $\times 40$); **B:** The ccRCC generally presents as round or polygonal, large distinctively transparent cells with thin-walled vascular network. (H&E staining, $\times 200$); **C:** The expression of PAX-8 in tumor tissue was detected by immunohistochemical technique, which indicates the tumor originated from the renal epithelial cells.

Reference

1. Bianchi M, Sun M, Jeldres C, Shariat SF, Trinh Q-D, Briganti A, et al. Distribution of metastatic sites in renal cell carcinoma: a population-based analysis. *Ann Oncol.* 2012; 23(4): 973-980.
2. Anwar SM, Kalbi DP, Upadhyaya A, Aqsa A, Mukherjee I. Scoping Renal Cell Carcinoma in the Rectal and Anal Canal: A Literature Review. *Cureus.* 2019; 11(12): e6330.
3. Dulskas A, Bagurskas P, Sinkevicius Z, Samalavicius NE. Sigmoid adenocarcinoma with metastases to the kidney: Report of a rare case and review of the literature. *Oncol Lett.* 2015; 10(2): 1191-1193.
4. Vedel PF, Wittendorf HE, Norus TP. Migration of clips to the colon after laparoscopic partial nephrectomy. *BMJ Case Rep.* 2017; 2017: bcr2016219040.